

Search for $f_J(2220)$ in radiative J/ψ decays

P. del Amo Sanchez,¹ J. P. Lees,¹ V. Poireau,¹ E. Prencipe,¹ V. Tisserand,¹ J. Garra Tico,² E. Grauges,² M. Martinelli,^{ab,3} A. Palano,^{ab,3} M. Pappagallo,^{ab,3} G. Eigen,⁴ B. Stugu,⁴ L. Sun,⁴ M. Battaglia,⁵ D. N. Brown,⁵ B. Hooberman,⁵ L. T. Kerth,⁵ Yu. G. Kolomensky,⁵ G. Lynch,⁵ I. L. Osipenko,⁵ T. Tanabe,⁵ C. M. Hawkes,⁶ A. T. Watson,⁶ H. Koch,⁷ T. Schroeder,⁷ D. J. Asgeirsson,⁸ C. Hearty,⁸ T. S. Mattison,⁸ J. A. McKenna,⁸ A. Khan,⁹ A. Randle-Conde,⁹ V. E. Blinov,¹⁰ A. R. Buzykaev,¹⁰ V. P. Druzhinin,¹⁰ V. B. Golubev,¹⁰ A. P. Onuchin,¹⁰ S. I. Serednyakov,¹⁰ Yu. I. Skovpen,¹⁰ E. P. Solodov,¹⁰ K. Yu. Todyshev,¹⁰ A. N. Yushkov,¹⁰ M. Bondioli,¹¹ S. Curry,¹¹ D. Kirkby,¹¹ A. J. Lankford,¹¹ M. Mandelkern,¹¹ E. C. Martin,¹¹ D. P. Stoker,¹¹ H. Atmacan,¹² J. W. Gary,¹² F. Liu,¹² O. Long,¹² G. M. Vitug,¹² C. Campagnari,¹³ T. M. Hong,¹³ D. Kovalskyi,¹³ J. D. Richman,¹³ A. M. Eisner,¹⁴ C. A. Heusch,¹⁴ J. Kroseberg,¹⁴ W. S. Lockman,¹⁴ A. J. Martinez,¹⁴ T. Schalk,¹⁴ B. A. Schumm,¹⁴ A. Seiden,¹⁴ L. O. Winstrom,¹⁴ C. H. Cheng,¹⁵ D. A. Doll,¹⁵ B. Echenard,¹⁵ D. G. Hitlin,¹⁵ P. Ongmongkolkul,¹⁵ F. C. Porter,¹⁵ A. Y. Rakitin,¹⁵ R. Andreassen,¹⁶ M. S. Dubrovin,¹⁶ G. Mancinelli,¹⁶ B. T. Meadows,¹⁶ M. D. Sokoloff,¹⁶ P. C. Bloom,¹⁷ W. T. Ford,¹⁷ A. Gaz,¹⁷ M. Nagel,¹⁷ U. Nauenberg,¹⁷ J. G. Smith,¹⁷ S. R. Wagner,¹⁷ R. Ayad,^{18,*} W. H. Toki,¹⁸ H. Jasper,¹⁹ T. M. Karbach,¹⁹ J. Merkel,¹⁹ A. Petzold,¹⁹ B. Spaan,¹⁹ K. Wacker,¹⁹ M. J. Kobel,²⁰ K. R. Schubert,²⁰ R. Schwierz,²⁰ D. Bernard,²¹ M. Verderi,²¹ P. J. Clark,²² S. Playfer,²² J. E. Watson,²² M. Andreotti,^{ab,23} D. Bettoni,^{a,23} C. Bozzi,^{a,23} R. Calabrese,^{ab,23} A. Cecchi,^{ab,23} G. Cibinetto,^{ab,23} E. Fioravanti,^{ab,23} P. Franchini,^{ab,23} E. Luppi,^{ab,23} M. Munerato,^{ab,23} M. Negrini,^{ab,23} A. Petrella,^{ab,23} L. Piemontese,^{a,23} R. Baldini-Ferrolì,²⁴ A. Calcaterra,²⁴ R. de Sangro,²⁴ G. Finocchiaro,²⁴ M. Nicolaci,²⁴ S. Pacetti,²⁴ P. Patteri,²⁴ I. M. Peruzzi,^{24,†} M. Piccolo,²⁴ M. Rama,²⁴ A. Zallo,²⁴ R. Contri,^{ab,25} E. Guido,^{ab,25} M. Lo Vetere,^{ab,25} M. R. Monge,^{ab,25} S. Passaggio,^{a,25} C. Patrignani,^{ab,25} E. Robutti,^{a,25} S. Tosi,^{ab,25} B. Bhuyan,²⁶ V. Prasad,²⁶ C. L. Lee,²⁷ M. Morii,²⁷ A. Adametz,²⁸ J. Marks,²⁸ U. Uwer,²⁸ F. U. Bernlochner,²⁹ M. Ebert,²⁹ H. M. Lacker,²⁹ T. Lueck,²⁹ A. Volk,²⁹ P. D. Dauncey,³⁰ M. Tibbetts,³⁰ P. K. Behera,³¹ U. Mallik,³¹ C. Chen,³² J. Cochran,³² H. B. Crawley,³² L. Dong,³² W. T. Meyer,³² S. Prell,³² E. I. Rosenberg,³² A. E. Rubin,³² Y. Y. Gao,³³ A. V. Gritsan,³³ Z. J. Guo,³³ N. Arnaud,³⁴ M. Davier,³⁴ D. Derkach,³⁴ J. Firmino da Costa,³⁴ G. Grosdidier,³⁴ F. Le Diberder,³⁴ A. M. Lutz,³⁴ B. Malaescu,³⁴ A. Perez,³⁴ P. Roudeau,³⁴ M. H. Schune,³⁴ J. Serrano,³⁴ V. Sordini,^{34,‡} A. Stocchi,³⁴ L. Wang,³⁴ G. Wormser,³⁴ D. J. Lange,³⁵ D. M. Wright,³⁵ I. Bingham,³⁶ C. A. Chavez,³⁶ J. P. Coleman,³⁶ J. R. Fry,³⁶ E. Gabathuler,³⁶ R. Gamet,³⁶ D. E. Hutchcroft,³⁶ D. J. Payne,³⁶ C. Touramanis,³⁶ A. J. Bevan,³⁷ F. Di Lodovico,³⁷ R. Sacco,³⁷ M. Sigamani,³⁷ G. Cowan,³⁸ S. Paramesvaran,³⁸ A. C. Wren,³⁸ D. N. Brown,³⁹ C. L. Davis,³⁹ A. G. Denig,⁴⁰ M. Fritsch,⁴⁰ W. Gradl,⁴⁰ A. Hafner,⁴⁰ K. E. Alwyn,⁴¹ D. Bailey,⁴¹ R. J. Barlow,⁴¹ G. Jackson,⁴¹ G. D. Lafferty,⁴¹ T. J. West,⁴¹ J. Anderson,⁴² R. Cenci,⁴² A. Jawahery,⁴² D. A. Roberts,⁴² G. Simi,⁴² J. M. Tuggle,⁴² C. Dallapiccola,⁴³ E. Salvati,⁴³ R. Cowan,⁴⁴ D. Dujmic,⁴⁴ P. H. Fisher,⁴⁴ G. Sciolla,⁴⁴ M. Zhao,⁴⁴ D. Lindemann,⁴⁵ P. M. Patel,⁴⁵ S. H. Robertson,⁴⁵ M. Schram,⁴⁵ P. Biassoni,^{ab,46} A. Lazzaro,^{ab,46} V. Lombardo,^{a,46} F. Palombo,^{ab,46} S. Stracka,^{ab,46} L. Cremaldi,⁴⁷ R. Godang,^{47,§} R. Kroeger,⁴⁷ P. Sonnek,⁴⁷ D. J. Summers,⁴⁷ X. Nguyen,⁴⁸ M. Simard,⁴⁸ P. Taras,⁴⁸ G. De Nardo,^{ab,49} D. Monorchio,^{ab,49} G. Onorato,^{ab,49} C. Sciacca,^{ab,49} G. Raven,⁵⁰ H. L. Snoek,⁵⁰ C. P. Jessop,⁵¹ K. J. Knoepfel,⁵¹ J. M. LoSecco,⁵¹ W. F. Wang,⁵¹ L. A. Corwin,⁵² K. Honscheid,⁵² R. Kass,⁵² J. P. Morris,⁵² N. L. Blount,⁵³ J. Brau,⁵³ R. Frey,⁵³ O. Igonkina,⁵³ J. A. Kolb,⁵³ R. Rahmat,⁵³ N. B. Sinev,⁵³ D. Strom,⁵³ J. Strube,⁵³ E. Torrence,⁵³ G. Castelli,^{ab,54} E. Feltres,^{ab,54} N. Gagliardi,^{ab,54} M. Margoni,^{ab,54} M. Morandin,^{a,54} M. Posocco,^{a,54} M. Rotondo,^{a,54} F. Simonetto,^{ab,54} R. Stroili,^{ab,54} E. Ben-Haim,⁵⁵ G. R. Bonneaud,⁵⁵ H. Briand,⁵⁵ G. Calderini,⁵⁵ J. Chauveau,⁵⁵ O. Hamon,⁵⁵ Ph. Leruste,⁵⁵ G. Marchiori,⁵⁵ J. Ocariz,⁵⁵ J. Prendki,⁵⁵ S. Sitt,⁵⁵ M. Biasini,^{ab,56} E. Manoni,^{ab,56} A. Rossi,^{ab,56} C. Angelini,^{ab,57} G. Batignani,^{ab,57} S. Bettarini,^{ab,57} M. Carpinelli,^{ab,57,¶} G. Casarosa,^{ab,57} A. Cervelli,^{ab,57} F. Forti,^{ab,57} M. A. Giorgi,^{ab,57} A. Lusiani,^{ac,57} N. Neri,^{ab,57} E. Paoloni,^{ab,57} G. Rizzo,^{ab,57} J. J. Walsh,^{a,57} D. Lopes Pegna,⁵⁸ C. Lu,⁵⁸ J. Olsen,⁵⁸ A. J. S. Smith,⁵⁸ A. V. Telnov,⁵⁸ F. Anulli,^{a,59} E. Baracchini,^{ab,59} G. Cavoto,^{a,59} R. Faccini,^{ab,59} F. Ferrarotto,^{a,59} F. Ferroni,^{ab,59} M. Gaspero,^{ab,59} L. Li Gioi,^{a,59} M. A. Mazzoni,^{a,59} G. Piredda,^{a,59} F. Renga,^{ab,59} T. Hartmann,⁶⁰ T. Leddig,⁶⁰ H. Schröder,⁶⁰ R. Waldi,⁶⁰ T. Adye,⁶¹ B. Franek,⁶¹ E. O. Olaiya,⁶¹ F. F. Wilson,⁶¹ S. Emery,⁶² G. Hamel de Monchenault,⁶² G. Vasseur,⁶² Ch. Yèche,⁶²

M. Zito,⁶² M. T. Allen,⁶³ D. Aston,⁶³ D. J. Bard,⁶³ R. Bartoldus,⁶³ J. F. Benitez,⁶³ C. Cartaro,⁶³ M. R. Convery,⁶³ J. Dorfan,⁶³ G. P. Dubois-Felsmann,⁶³ W. Dunwoodie,⁶³ R. C. Field,⁶³ M. Franco Sevilla,⁶³ B. G. Fulsom,⁶³ A. M. Gabareen,⁶³ M. T. Graham,⁶³ P. Grenier,⁶³ C. Hast,⁶³ W. R. Innes,⁶³ M. H. Kelsey,⁶³ H. Kim,⁶³ P. Kim,⁶³ M. L. Kocian,⁶³ D. W. G. S. Leith,⁶³ S. Li,⁶³ B. Lindquist,⁶³ S. Luitz,⁶³ V. Luth,⁶³ H. L. Lynch,⁶³ D. B. MacFarlane,⁶³ H. Marsiske,⁶³ D. R. Muller,⁶³ H. Neal,⁶³ S. Nelson,⁶³ C. P. O'Grady,⁶³ I. Ofte,⁶³ M. Perl,⁶³ T. Pulliam,⁶³ B. N. Ratcliff,⁶³ A. Roodman,⁶³ A. A. Salnikov,⁶³ V. Santoro,⁶³ R. H. Schindler,⁶³ J. Schwiening,⁶³ A. Snyder,⁶³ D. Su,⁶³ M. K. Sullivan,⁶³ S. Sun,⁶³ K. Suzuki,⁶³ J. M. Thompson,⁶³ J. Va'vra,⁶³ A. P. Wagner,⁶³ M. Weaver,⁶³ C. A. West,⁶³ W. J. Wisniewski,⁶³ M. Wittgen,⁶³ D. H. Wright,⁶³ H. W. Wulsin,⁶³ A. K. Yarritu,⁶³ C. C. Young,⁶³ V. Ziegler,⁶³ X. R. Chen,⁶⁴ W. Park,⁶⁴ M. V. Purohit,⁶⁴ R. M. White,⁶⁴ J. R. Wilson,⁶⁴ S. J. Sekula,⁶⁵ M. Bellis,⁶⁶ P. R. Burchat,⁶⁶ A. J. Edwards,⁶⁶ T. S. Miyashita,⁶⁶ S. Ahmed,⁶⁷ M. S. Alam,⁶⁷ J. A. Ernst,⁶⁷ B. Pan,⁶⁷ M. A. Saeed,⁶⁷ S. B. Zain,⁶⁷ N. Guttman,⁶⁸ A. Soffer,⁶⁸ P. Lund,⁶⁹ S. M. Spanier,⁶⁹ R. Eckmann,⁷⁰ J. L. Ritchie,⁷⁰ A. M. Ruland,⁷⁰ C. J. Schilling,⁷⁰ R. F. Schwitters,⁷⁰ B. C. Wray,⁷⁰ J. M. Izen,⁷¹ X. C. Lou,⁷¹ F. Bianchi^{ab,72} D. Gamba^{ab,72} M. Pelliccioni^{ab,72} M. Bomben^{ab,73} L. Lanceri^{ab,73} L. Vitale^{ab,73} N. Lopez-March,⁷⁴ F. Martinez-Vidal,⁷⁴ D. A. Milanes,⁷⁴ A. Oyanguren,⁷⁴ J. Albert,⁷⁵ Sw. Banerjee,⁷⁵ H. H. F. Choi,⁷⁵ K. Hamano,⁷⁵ G. J. King,⁷⁵ R. Kowalewski,⁷⁵ M. J. Lewczuk,⁷⁵ I. M. Nugent,⁷⁵ J. M. Roney,⁷⁵ R. J. Sobie,⁷⁵ T. J. Gershon,⁷⁶ P. F. Harrison,⁷⁶ T. E. Latham,⁷⁶ E. M. T. Puccio,⁷⁶ H. R. Band,⁷⁷ S. Dasu,⁷⁷ K. T. Flood,⁷⁷ Y. Pan,⁷⁷ R. Prepost,⁷⁷ C. O. Vuosalo,⁷⁷ and S. L. Wu⁷⁷

(The BABAR Collaboration)

¹Laboratoire d'Annecy-le-Vieux de Physique des Particules (LAPP),
Université de Savoie, CNRS/IN2P3, F-74941 Annecy-Le-Vieux, France

²Universitat de Barcelona, Facultat de Física, Departament ECM, E-08028 Barcelona, Spain

³INFN Sezione di Bari^a; Dipartimento di Fisica, Università di Bari^b, I-70126 Bari, Italy

⁴University of Bergen, Institute of Physics, N-5007 Bergen, Norway

⁵Lawrence Berkeley National Laboratory and University of California, Berkeley, California 94720, USA

⁶University of Birmingham, Birmingham, B15 2TT, United Kingdom

⁷Ruhr Universität Bochum, Institut für Experimentalphysik 1, D-44780 Bochum, Germany

⁸University of British Columbia, Vancouver, British Columbia, Canada V6T 1Z1

⁹Brunel University, Uxbridge, Middlesex UB8 3PH, United Kingdom

¹⁰Budker Institute of Nuclear Physics, Novosibirsk 630090, Russia

¹¹University of California at Irvine, Irvine, California 92697, USA

¹²University of California at Riverside, Riverside, California 92521, USA

¹³University of California at Santa Barbara, Santa Barbara, California 93106, USA

¹⁴University of California at Santa Cruz, Institute for Particle Physics, Santa Cruz, California 95064, USA

¹⁵California Institute of Technology, Pasadena, California 91125, USA

¹⁶University of Cincinnati, Cincinnati, Ohio 45221, USA

¹⁷University of Colorado, Boulder, Colorado 80309, USA

¹⁸Colorado State University, Fort Collins, Colorado 80523, USA

¹⁹Technische Universität Dortmund, Fakultät Physik, D-44221 Dortmund, Germany

²⁰Technische Universität Dresden, Institut für Kern- und Teilchenphysik, D-01062 Dresden, Germany

²¹Laboratoire Leprince-Ringuet, CNRS/IN2P3, Ecole Polytechnique, F-91128 Palaiseau, France

²²University of Edinburgh, Edinburgh EH9 3JZ, United Kingdom

²³INFN Sezione di Ferrara^a; Dipartimento di Fisica, Università di Ferrara^b, I-44100 Ferrara, Italy

²⁴INFN Laboratori Nazionali di Frascati, I-00044 Frascati, Italy

²⁵INFN Sezione di Genova^a; Dipartimento di Fisica, Università di Genova^b, I-16146 Genova, Italy

²⁶Indian Institute of Technology Guwahati, Guwahati, Assam, 781 039, India

²⁷Harvard University, Cambridge, Massachusetts 02138, USA

²⁸Universität Heidelberg, Physikalisches Institut, Philosophenweg 12, D-69120 Heidelberg, Germany

²⁹Humboldt-Universität zu Berlin, Institut für Physik, Newtonstr. 15, D-12489 Berlin, Germany

³⁰Imperial College London, London, SW7 2AZ, United Kingdom

³¹University of Iowa, Iowa City, Iowa 52242, USA

³²Iowa State University, Ames, Iowa 50011-3160, USA

³³Johns Hopkins University, Baltimore, Maryland 21218, USA

³⁴Laboratoire de l'Accélérateur Linéaire, IN2P3/CNRS et Université Paris-Sud 11,
Centre Scientifique d'Orsay, B. P. 34, F-91898 Orsay Cedex, France

³⁵Lawrence Livermore National Laboratory, Livermore, California 94550, USA

³⁶University of Liverpool, Liverpool L69 7ZE, United Kingdom

³⁷Queen Mary, University of London, London, E1 4NS, United Kingdom

³⁸University of London, Royal Holloway and Bedford New College, Egham, Surrey TW20 0EX, United Kingdom

³⁹University of Louisville, Louisville, Kentucky 40292, USA

- ⁴⁰ Johannes Gutenberg-Universität Mainz, Institut für Kernphysik, D-55099 Mainz, Germany
⁴¹ University of Manchester, Manchester M13 9PL, United Kingdom
⁴² University of Maryland, College Park, Maryland 20742, USA
⁴³ University of Massachusetts, Amherst, Massachusetts 01003, USA
⁴⁴ Massachusetts Institute of Technology, Laboratory for Nuclear Science, Cambridge, Massachusetts 02139, USA
⁴⁵ McGill University, Montréal, Québec, Canada H3A 2T8
⁴⁶ INFN Sezione di Milano^a; Dipartimento di Fisica, Università di Milano^b, I-20133 Milano, Italy
⁴⁷ University of Mississippi, University, Mississippi 38677, USA
⁴⁸ Université de Montréal, Physique des Particules, Montréal, Québec, Canada H3C 3J7
⁴⁹ INFN Sezione di Napoli^a; Dipartimento di Scienze Fisiche, Università di Napoli Federico II^b, I-80126 Napoli, Italy
⁵⁰ NIKHEF, National Institute for Nuclear Physics and High Energy Physics, NL-1009 DB Amsterdam, The Netherlands
⁵¹ University of Notre Dame, Notre Dame, Indiana 46556, USA
⁵² Ohio State University, Columbus, Ohio 43210, USA
⁵³ University of Oregon, Eugene, Oregon 97403, USA
⁵⁴ INFN Sezione di Padova^a; Dipartimento di Fisica, Università di Padova^b, I-35131 Padova, Italy
⁵⁵ Laboratoire de Physique Nucléaire et de Hautes Energies, IN2P3/CNRS, Université Pierre et Marie Curie-Paris6, Université Denis Diderot-Paris7, F-75252 Paris, France
⁵⁶ INFN Sezione di Perugia^a; Dipartimento di Fisica, Università di Perugia^b, I-06100 Perugia, Italy
⁵⁷ INFN Sezione di Pisa^a; Dipartimento di Fisica, Università di Pisa^b; Scuola Normale Superiore di Pisa^c, I-56127 Pisa, Italy
⁵⁸ Princeton University, Princeton, New Jersey 08544, USA
⁵⁹ INFN Sezione di Roma^a; Dipartimento di Fisica, Università di Roma La Sapienza^b, I-00185 Roma, Italy
⁶⁰ Universität Rostock, D-18051 Rostock, Germany
⁶¹ Rutherford Appleton Laboratory, Chilton, Didcot, Oxon, OX11 0QX, United Kingdom
⁶² CEA, Irfu, SPP, Centre de Saclay, F-91191 Gif-sur-Yvette, France
⁶³ SLAC National Accelerator Laboratory, Stanford, California 94309 USA
⁶⁴ University of South Carolina, Columbia, South Carolina 29208, USA
⁶⁵ Southern Methodist University, Dallas, Texas 75275, USA
⁶⁶ Stanford University, Stanford, California 94305-4060, USA
⁶⁷ State University of New York, Albany, New York 12222, USA
⁶⁸ Tel Aviv University, School of Physics and Astronomy, Tel Aviv, 69978, Israel
⁶⁹ University of Tennessee, Knoxville, Tennessee 37996, USA
⁷⁰ University of Texas at Austin, Austin, Texas 78712, USA
⁷¹ University of Texas at Dallas, Richardson, Texas 75083, USA
⁷² INFN Sezione di Torino^a; Dipartimento di Fisica Sperimentale, Università di Torino^b, I-10125 Torino, Italy
⁷³ INFN Sezione di Trieste^a; Dipartimento di Fisica, Università di Trieste^b, I-34127 Trieste, Italy
⁷⁴ IFIC, Universitat de Valencia-CSIC, E-46071 Valencia, Spain
⁷⁵ University of Victoria, Victoria, British Columbia, Canada V8W 3P6
⁷⁶ Department of Physics, University of Warwick, Coventry CV4 7AL, United Kingdom
⁷⁷ University of Wisconsin, Madison, Wisconsin 53706, USA

(Dated: September 10, 2010)

We present a search for $f_J(2220)$ production in radiative $J/\psi \rightarrow \gamma f_J(2220)$ decays using 460 fb⁻¹ of data collected with the BABAR detector at the SLAC PEP-II e^+e^- collider. The $f_J(2220)$ is searched for in the decays to K^+K^- and $K_S^0K_S^0$. No evidence of this resonance is observed, and 90% confidence level upper limits on the product of the branching fractions for $J/\psi \rightarrow \gamma f_J(2220)$ and $f_J(2220) \rightarrow K^+K^- (K_S^0K_S^0)$ as a function of spin and helicity are set at the level of 10^{-5} , below the central values reported by the Mark III experiment.

PACS numbers: 12.39.Mk, 13.20.Gd

Evidence for the $f_J(2220)$, a narrow resonance with a mass around 2.2 GeV/ c^2 also known as $\xi(2230)$, was first presented by the Mark III Collaboration [1]. The $f_J(2220)$ was seen as a narrow signal above a broad enhancement in both $J/\psi \rightarrow \gamma f_J(2220)$, $f_J(2220) \rightarrow K^+K^-$ and $J/\psi \rightarrow \gamma f_J(2220)$, $f_J(2220) \rightarrow K_S^0K_S^0$ decays. The charged and neutral product branching fractions (PBF) were measured to be $(4.2_{-1.4}^{+1.7} \pm 0.8) \times 10^{-5}$

and $(3.1_{-1.3}^{+1.6} \pm 0.7) \times 10^{-5}$ with significance of 3.6 and 4.7 standard deviations, respectively. The BES Collaboration has also subsequently reported evidence in radiative J/ψ decays at a comparable level of significance [2]. They reported PBF of $(3.3_{-1.3}^{+1.6} \pm 1.2) \times 10^{-5}$ and $(2.7_{-0.9}^{+1.1} \pm 0.8) \times 10^{-5}$ for the K^+K^- and $K_S^0K_S^0$ channels. Indications of similar structure produced in π^-p and K^-p collisions have been seen [3–5], while searches

for direct formation in $p\bar{p}$ collisions [6, 7] or two-photon processes [8, 9] were inconclusive.

The unexpectedly narrow width of the $f_J(2220)$, approximately 20 MeV, triggered speculation about its nature. Besides the early hypothesis of a “light Higgs” [10], conjectures range from a multiquark state, to a hybrid resonance, a $\Lambda\bar{\Lambda}$ bound state, a high-spin $s\bar{s}$ state, or a glueball [11]. Intriguingly, lattice QCD calculations predict a mass for the ground state tensor 2^{++} glueball close to 2.2 GeV/ c^2 [12, 13].

We report herein a search for the $f_J(2220)$ in radiative J/ψ decays, with the J/ψ produced via initial state radiation (ISR) in e^+e^- collisions recorded at PEP-II. The emission of ISR allows the study of resonance production over a wide range of e^+e^- center-of-mass (CM) energies [14]. The data sample used in this analysis consists of 425 fb $^{-1}$ recorded at $\sqrt{s} = 10.54$ GeV and 35 fb $^{-1}$ recorded 40 MeV below this energy. With a luminosity-weighted cross section for J/ψ production of 35.7 pb, this dataset contains 16.4 ± 0.3 million directly-produced J/ψ decays.

The BABAR detector is described in detail elsewhere [15]. Charged particle momenta are measured in a tracking system consisting of a five-layer double-sided silicon vertex detector (SVT), and a 40-layer central drift chamber (DCH), immersed in a 1.5-T axial magnetic field. Photon and electron energies are measured in a CsI(Tl) electromagnetic calorimeter (EMC). Charged particle identification (PID) is performed using an internally reflecting ring-imaging Cherenkov detector and the energy loss dE/dx , measured by the SVT and DCH.

Detector acceptance is studied using Monte Carlo (MC) simulation based on GEANT4 [16]. Multiple photon emission from the initial-state charged particles is implemented using a structure function technique [17, 18]. The $f_J(2220)$ resonance is modeled by a non-relativistic Breit-Wigner function with a mass of 2.231 GeV/ c^2 and a width of 23 MeV [19]. Several hypotheses for the spin and helicity of the $f_J(2220)$ are considered: spin $J = 0$, and spin $J = 2$ with pure helicity ± 2 , ± 1 or 0. The hypothesis $J = 4$ is strongly disfavored by lattice QCD calculations [20].

The $J/\psi \rightarrow \gamma K^+ K^-$ decay is reconstructed by combining two oppositely charged tracks, identified as kaons, with a photon candidate. Events containing a π^0 candidate, defined as a pair of photons of energy larger than 50 MeV [21] having an invariant mass in the range 115 – 155 MeV/ c^2 , are discarded. The contamination of $J/\psi \rightarrow K^{*\pm}(892)(K^\pm\pi^0)K^\mp$, in which the π^0 is not reconstructed, is further reduced by rejecting J/ψ candidates having a kaon with a momentum larger than 1.35 GeV/ c in the J/ψ CM frame.

The $J/\psi \rightarrow \gamma K_S^0 K_S^0$ channel, examined in $J/\psi \rightarrow \gamma\pi^+\pi^-\pi^+\pi^-$, is reconstructed using events containing a photon and four charged tracks. Neutral kaon candidates are reconstructed from $K_S^0 \rightarrow \pi^+\pi^-$, combining a pair of oppositely charged tracks identified as pions, with an in-

variant mass in the range $|M_{\pi^+\pi^-} - M_{K_S}| < 15$ MeV/ c^2 . To improve the signal purity, the angle in the transverse plane between the momentum and the flight direction of each kaon is required to be less than 0.1 rad. No π^0 veto is applied, as the $J/\psi \rightarrow K_S^0 K_S^0 \pi^0$ decay is forbidden by C -parity conservation and the overall π^0 contamination is negligible.

Events with additional charged tracks are rejected. The photon emitted by the J/ψ is also required to have an energy larger than 300 MeV to suppress background from additional ISR photons or noise from the calorimeter. Finally, the helicity angle of each kaon, ζ_K , must satisfy $|\cos \zeta_K| < 0.7$.




FIG. 1: The distribution of M_{rec}^2 , the square of the recoiling mass against the $J/\psi \rightarrow \gamma K^+ K^-$ (a) and $J/\psi \rightarrow \gamma K_S^0 K_S^0$ (b) candidates, after all other selection criteria are applied for events in which the ISR photon is detected (open circle) or undetected (solid circle).

Radiative $e^+e^- \rightarrow \gamma_{ISR} J/\psi$ events are then identified. Clusters in the EMC not associated with charged-particle tracks and having energy larger than 1 GeV are taken as ISR photon candidates. Events in which the ISR photon falls within the detector acceptance are selected by demanding an angle between the J/ψ candidate and the ISR photon in the CM frame larger than 3.12 (3.10) rad for the charged (neutral) mode. In the opposite case, the square of the mass recoiling against the J/ψ is required to lie between -2.0 GeV $^2/c^4$ (-2.0 GeV $^2/c^4$) and 2.0 GeV $^2/c^4$ (5.0 GeV $^2/c^4$) for $J/\psi \rightarrow \gamma K^+ K^-$ ($K_S^0 K_S^0$) candidates. In both cases, no additional photons with energy exceeding 300 MeV can be present. For the charged mode, the cosine of the polar angle of the photon emitted by the J/ψ is required to be less than 0.8, and, for events where the ISR photon is undetected, that of each kaon must be less than 0.9. The distribution of the recoiling mass squared after applying all other cuts is displayed in Fig. 1 for combinations having a mass in the range $2.8 < m_{\gamma KK} < 3.4$ GeV/ c^2 . Clear peaks corresponding to ISR events are visible.

The resulting $\gamma K^+ K^-$ and $\gamma K_S^0 K_S^0$ mass distributions are displayed in Fig. 2. A large J/ψ signal over a smooth background is observed for both channels. This background, hereafter referred to as inclusive, arises mainly

from partially-reconstructed $J/\psi \rightarrow KK + X$ decays and $e^+e^- \rightarrow q\bar{q}\gamma_{ISR}$ ($q = u, d, s, c$) production. Its level in the J/ψ region is determined by fitting the data with a Gaussian and a second (first) order polynomial for the charged (neutral) mode. The J/ψ candidates are then fitted, constraining their mass to the world-average value [19] and requiring a common vertex for the decay products. A mass constraint on both K_S^0 candidates is also imposed for the neutral channel. Combinations having a fit probability larger than 0.01 are retained to form the final sample. The corresponding inclusive background is evaluated by correcting the values extrapolated from the unconstrained mass spectra for the efficiency of the fit fit probability cut.




FIG. 2: The $\gamma K^+ K^-$ (a) and $\gamma K_S^0 K_S^0$ (b) mass spectra after all selection criteria are applied. The points represent data, and the plain histograms show combinations having fit probability larger than 0.01. The estimated inclusive background in the final sample is shown as a filled histogram.

The fitted $K^+ K^-$ and $K_S^0 K_S^0$ mass spectra are shown in Fig. 3, together with the contribution of various J/ψ decays and the inclusive background. The shape of the inclusive background is modeled using sideband data taken from the unconstrained mass spectra in the ranges $2.7 < m_{\gamma KK} < 2.9 \text{ GeV}/c^2$ and $3.2 < m_{\gamma KK} < 3.4 \text{ GeV}/c^2$. The contributions of the $J/\psi \rightarrow \gamma f_2'(1525)$, $J/\psi \rightarrow \gamma f_0(1710)$, and $J/\psi \rightarrow K^{\pm} K^{\mp}$ channels are estimated from MC simulation using world-average branching fractions [19]. Contamination from $J/\psi \rightarrow K^{*\pm} K^{\mp}$ decays is found to be negligible. The $f_2'(1525) \rightarrow K^+ K^-$ and $f_2'(1525) \rightarrow K_S^0 K_S^0$ decays are modeled using helicity amplitude ratios $x^2 = 1.0$ and $y^2 = 0.44$ [22]. No interference between the $f_0(1710)$ and the inclusive background is considered. The sum of these components accounts for most of the data in the region below $2 \text{ GeV}/c^2$ and reproduces well the contribution of $\phi(1020)$ mesons. The excess seen around $1.25 - 1.30 \text{ GeV}/c^2$ in the charged mode is likely due to $J/\psi \rightarrow \rho^0 \pi^0, \rho^0 \rightarrow \pi^+ \pi^-$ decays, where both charged pions are misidentified as kaons, and a photon from the π^0 decay goes undetected. The data above $2 \text{ GeV}/c^2$ are dominated by partially-reconstructed J/ψ decays.

The number of signal events is determined using an unbinned maximum likelihood fit in the range $1.9 \text{ GeV}/c^2 <$




FIG. 3: The fitted $K^+ K^-$ (a) and $K_S^0 K_S^0$ (b) mass spectra. The expected contributions of the inclusive background (plain histogram), $J/\psi \rightarrow \gamma f_2'(1525)$ (cross-hatched histogram), $J/\psi \rightarrow \gamma f_0(1710)$ (hatched histogram) are also shown. The results of the fits are displayed in the inserts.

$m_{KK} < 2.6 \text{ GeV}/c^2$. The signal is described by a Breit-Wigner distribution convolved with a Gaussian resolution function, while the background is modeled by a second order Chebychev polynomial. The mass and width of the resonance are fixed to $2.231 \text{ GeV}/c^2$ and 23 MeV , respectively. The Gaussian resolution, taken from MC simulations, is set to $8 \text{ MeV}/c^2$ ($6 \text{ MeV}/c^2$) for the $K^+ K^-$ ($K_S^0 K_S^0$) channel. We have checked on a number of independent control samples that the two-body invariant mass resolution is well reproduced by the MC over the whole invariant mass range studied in this paper. The results of the fits are displayed in Fig. 3. No evidence of a $f_J(2220)$ signal is observed.

The largest sources of systematic uncertainty arise from the parametrization of the signal and background shapes. An uncertainty of 0.2 events arises from fixing the mass, width and resolution of the signal in each channel. This contribution is estimated by varying each parameter by $\pm 1\sigma$ in the fitting procedure. Similarly, the uncertainty due to the background parametrization, evaluated to be 1.4 (0.6) events for the $K^+ K^-$ ($K_S^0 K_S^0$) mode, is assessed by repeating the fit with a third order Chebychev polynomial. Multiplicative systematic uncer-

TABLE I: The efficiency, the PBF of the decays $J/\psi \rightarrow \gamma f_J(2220)$, $f_J(2220) \rightarrow K^+K^-$ and $J/\psi \rightarrow \gamma f_J(2220)$, $f_J(2220) \rightarrow K_S^0 K_S^0$ and corresponding 90% confidence level upper limit (UL) as a function of the spin J and helicity h assumed for the $f_J(2220)$. The number of $f_J(2220) \rightarrow K^+K^- (K_S^0 K_S^0)$ events determined from the fit is $1.0^{+8.9}_{-7.9} \pm 1.5$ ($-0.8^{+2.1}_{-1.2} \pm 0.6$). The first uncertainty is statistical and the second systematic.

Spin / helicity hypothesis	Efficiency (%)	PBF ($\times 10^{-5}$)	UL ($\times 10^{-5}$)
$f_J(2220) \rightarrow K^+K^-$			
$J = 0$	5.15 ± 0.03	$0.12^{+1.05}_{-0.94} \pm 0.17$	< 1.9
$J = 2 / h = 0$	2.74 ± 0.04	$0.22^{+1.97}_{-1.76} \pm 0.33$	< 3.6
$J = 2 / h = \pm 1$	5.22 ± 0.05	$0.12^{+1.03}_{-0.93} \pm 0.17$	< 1.9
$J = 2 / h = \pm 2$	6.69 ± 0.05	$0.09^{+0.81}_{-0.72} \pm 0.13$	< 1.5
$f_J(2220) \rightarrow K_S^0 K_S^0$			
$J = 0$	1.32 ± 0.01	$-0.39^{+0.96}_{-0.56} \pm 0.28$	< 1.7
$J = 2 / h = 0$	0.74 ± 0.01	$-0.69^{+0.71}_{-1.00} \pm 0.49$	< 2.9
$J = 2 / h = \pm 1$	1.39 ± 0.02	$-0.37^{+0.92}_{-0.54} \pm 0.26$	< 1.6
$J = 2 / h = \pm 2$	1.75 ± 0.02	$-0.29^{+0.73}_{-0.42} \pm 0.21$	< 1.2

tainties on the charged (neutral) PBF include the selection procedure [4.0% (2.2%)], the determination of the number of J/ψ mesons [3.0% (3.0%)], the trigger efficiencies [3.1% (3.5%)], the track and neutral cluster reconstruction [1.9% (3.3%)], the particle identification [1.4% (-)], and the MC statistics [1.0% (1.4%)].

The $J/\psi \rightarrow \gamma f_J(2220)$, $f_J(2220) \rightarrow K^+K^-$ and $J/\psi \rightarrow \gamma f_J(2220)$, $f_J(2220) \rightarrow K_S^0 K_S^0$ PBF are given in Table I as a function of the spin and helicity assumed for the $f_J(2220)$. The efficiencies are determined from the corresponding MC and include the $K_S^0 \rightarrow \pi^+ \pi^-$ branching fraction as well as corrections for particle identification, photon detection and K_S^0 reconstruction. The 90% confidence level (CL) Bayesian upper limits, based on priors uniform in branching fraction and including systematic uncertainties, are also shown.

In conclusion, no evidence is observed for the $f_J(2220)$ in radiative J/ψ decay in ISR events produced in e^+e^- collisions at $\sqrt{s} = m_{\Upsilon(4S)}$. For all hypotheses of spin and helicity, the 90% CL upper limits on the $J/\psi \rightarrow \gamma f_J(2220)$, $f_J(2220) \rightarrow K^+K^-$ and $J/\psi \rightarrow \gamma f_J(2220)$, $f_J(2220) \rightarrow K_S^0 K_S^0$ PBF are below the central values reported by Mark III. Only one hypothesis of spin and helicity ($J = 2$ and $h = 0$) is compatible with the BES results for both final states, while all other possibilities are clearly excluded.

We are grateful for the excellent luminosity and machine conditions provided by our PEP-II colleagues, and for the substantial dedicated effort from the computing organizations that support BABAR. The collaborating institutions wish to thank SLAC for its support and kind hospitality. This work is supported by DOE and NSF (USA), NSERC (Canada), CEA and CNRS-IN2P3

(France), BMBF and DFG (Germany), INFN (Italy), FOM (The Netherlands), NFR (Norway), MES (Russia), MICIIN (Spain), STFC (United Kingdom). Individuals have received support from the Marie Curie EIF (European Union), the A. P. Sloan Foundation (USA) and the Binational Science Foundation (USA-Israel).

* Now at Temple University, Philadelphia, Pennsylvania 19122, USA

† Also with Università di Perugia, Dipartimento di Fisica, Perugia, Italy

‡ Also with Università di Roma La Sapienza, I-00185 Roma, Italy

§ Now at University of South Alabama, Mobile, Alabama 36688, USA

¶ Also with Università di Sassari, Sassari, Italy

- [1] R. M. Baltrusaitis *et al.* (Mark III Collab.), Phys. Rev. Lett. **56**, 107 (1986).
- [2] J. Z. Bai *et al.* (BES Collab.), Phys. Rev. Lett. **76**, 3502 (1996).
- [3] B. V. Bolonkin *et al.*, Yad. Fiz. **46** 799 (1987) [Nucl. Phys. B **309** 426 (1988)].
- [4] D. Aston *et al.*, Phys. Lett. B **215** 199 (1988).
- [5] D. Alde *et al.*, Phys. Lett. B **177** 120 (1986).
- [6] C. Amsler *et al.* (Crystal Ball Collab.), Phys. Lett. B **520**, 175 (2001).
- [7] C. Evangelista *et al.* (JETSET Collab.), Phys. Rev. D **57**, 5370 (1998) and references therein.
- [8] K. Benslama *et al.* (CLEO Collab.), Phys. Rev. D **66**, 077101 (2002).
- [9] M. Acciarri *et al.* (L3 Collab.), Phys. Lett. B **501**, 173 (2001).
- [10] R. M. Barnett, G. Senjanovic and D. Wyler, Phys. Rev. D **30**, 1529 (1984) and references therein.
- [11] M. S. Chanowitz and S. R. Sharpe, Phys. Lett. B **132**, 413 (1983); K.-T. Chao, Commun. Theor. Phys. **3**, 757 (1984); S. Pakvasa, M. Suzuki, and S. F. Tuan, Phys. Lett. B **145**, 135 (1984); M. P. Shatz, Phys. Lett. B **138**, 209 (1984); A. Le Yaouanc *et al.*, Z. Phys. C **28**, 309 (1985); S. Godfrey, R. Kokoski and N. Isgur, Phys. Lett. B **141**, 439 (1984); B. F. L. Ward, Phys. Rev. D **31**, 2849 (1985) [Erratum-ibid. D **32**, 1260 (1985)]. S. Ono, Phys. Rev. D **35**, 944 (1987); K.-T. Chao, Phys. Rev. Lett. **60**, 2579 (1988);
- [12] J.-X. Chen and J.-C. Su, Phys. Rev. D **69**, 076003 (2004).
- [13] C. J. Morningstar and M. J. Peardon, Phys. Rev. D **56**, 4043 (1997).
- [14] See for example M. Benayoun *et al.*, Mod. Phys. Lett. A **14**, 2605 (1999).
- [15] B. Aubert *et al.* (BABAR Collab.), Nucl. Instrum. Meth. A **479**, 1 (2002).
- [16] S. Agostinelli *et al.* (GEANT4 Collab.), Nucl. Instrum. Meth. A **506**, 250 (2003).
- [17] A. B. Arbuzov *et al.*, J. High Energy Phys. **9710**, 001 (1997).
- [18] M. Caffo, H. Czyż, and E. Remiddi, Nuovo Cim. **A110**, 515 (1997); Phys. Lett. **B327**, 369 (1994).
- [19] C. Amsler *et al.* (Particle Data Group), Phys. Lett. B **667**, 1 (2008) and 2009 update.

- [20] D. Q. Liu and J. M. Wu, Mod. Phys. Lett. A **17**, 1419 (2002).
- [21] All kinematic quantities are defined in the laboratory frame unless another frame is specified.
- [22] J. Z. Bai *et al.* (BES Collab.), Phys. Rev. D **68**, 052003 (2003).